



Scopus

Document details

[Back to results](#) | 1 of 1
[Export](#)
[Download](#)
[Print](#)
[E-mail](#)
[Save to PDF](#)
[Add to List](#)
[More...](#)

Journal of Pure and Applied Microbiology
Volume 8, Issue SPEC. ISS. 1, May 2014, Pages 769-776

Identification of microbes in inoculant for agarwood (Gaharu) formation using molecular biological techniques (Article)

Nuge, T.^a, Hashim, Y.Z.H.-Y.^{ab}, Amid, A.^{ab}, Jaswir, I.^{ab}, Jamal, P.^{ab}, Salleh, H.M.^{ab}  

^aDepartment of Biotechnology Engineering, Faculty of Engineering, International Islamic University Malaysia, Malaysia

^bInternational Institute for Halal Research and Training (INHART), International Islamic University Malaysia, Malaysia

Abstract

[View references \(22\)](#)

The distinctive fragrance and pleasing odor of the agarwood came from the resinous materials impregnated in the phloem of the tree as a response to infection. Artificial inoculants have been developed by "trial and error" methods to boost the resin formation. These artificial inoculants have been used widely in this region despite no strong scientific data and that they contribute to most of the cost for agarwood resin production. In collaboration with an agarwood industry player, a research work was initiated to isolate and identify the microorganisms present in an inoculant that has been proven effective in boosting the agarwood resin formation. Fifteen clones were constructed by extracting and amplifying the 16S rDNA and ITS sequences from the pure cultures isolated from the inoculant. Three genera of microorganisms (*Bacillus*, *Paenibacillus*, *Monascus*) have been successfully identified. Two bacteria clones were phylogenetically affiliated with *Paenibacillus* sp. and four clones were associated with *Bacillus* sp. as well as three fungi clones were associated with *Monascus* sp. All the identified species are spore-forming microorganisms. The genera of *Bacillus* and *Paenibacillus* were able to utilize carbohydrate and protein from the tree as their energy sources. These two genera may be crucial to boost resin formation in agarwood.

Author keywords

16S rDNA Agarwood Inoculant ITS Phylogeny

ISSN: 09737510

Source Type: Journal

Original language: English

Document Type: Article

Publisher: Journal of Pure and Applied Microbiology

References (22)

[View in search results format >](#)

☐ All [Export](#) [Print](#) [E-mail](#) [Save to PDF](#) [Create bibliography](#)

☐ 1 Fui, L.H., Mamat, M.P., Shyun, C.Y.
Production use and trade of Gaharu in Peninsular Malaysia
International Economic Conference on Trade & Industry: Georgetown, Penang, 2007

☐ 2 Barden, A., AwangAnak, N., Mulliken, T., Song, M.
Heart of the Matter: Agarwood Use and Trade, and CITES implementation for *Aquilaria malaccensis*
(2000) *TRAFFIC International Report*

Metrics

0 Citations in Scopus

0 Field-Weighted
Citation Impact



PlumX Metrics 

Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 0 documents

Inform me when this document
is cited in Scopus:

[Set citation alert >](#)

[Set citation feed >](#)

Related documents

No More Green Thumbs!

Bland, J.A.
(1977) *American Biology Teacher*

First report of leaf spot of
dracaena reflexa caused by
burkholderia gladioli worldwide

Romberg, M.K. , Griffin, R.L. ,
Murugan, S.
(2010) *Plant Disease*

Comparison of the characteristics
of *Monascus* Strains in the solid-
state fermentation system

Liang, S. , Ni, L.
(2015) *Journal of Chinese
Institute of Food Science and
Technology*

View all related documents based
on references

Find more related documents in
Scopus based on:

[Authors >](#) [Keywords >](#)